



# Resin tapping influence on maritime pine growth depends on tree age and stand characteristics

Mikael Moura<sup>1</sup> · Filipe Campelo<sup>1</sup> · Cristina Nabais<sup>1</sup> · Núria Garcia-Forner<sup>1</sup>

Received: 22 April 2022 / Revised: 9 February 2023 / Accepted: 3 April 2023  
© The Author(s) 2023

## Abstract

Resin is a renewable forest resource that can increase the economic value of some forests in rural areas. Resin production is associated with climatic conditions. However, its impact on trees' growth remains unclear. Here, we studied radial growth in six Portuguese *Pinus pinaster* forests that had been resin tapped for 5–7 years along a latitudinal and climatic gradient to understand whether resin tapping affects tree growth, and how it is affected by climate, stand and tree traits. Tree-ring width (*TRW*) on tapped and untapped trunk sides was compared before and during the tapping period. Tree-ring width decreased in the three youngest populations (< 30 years), with no changes in older populations (> 40 years), while *TRW* increased after resin harvesting began in the oldest stand (> 55 years). Annual resin-tapping impact (*RTI*), calculated as the ratio between *TRW* during tapping years and the 5-year average *TRW* before tapping, was below and above 1 in the younger and older stands, respectively. Among stand characteristics and across sites, *RTI* was negatively correlated with tree competition and positively correlated with *TRW*, cambial age, and tree height. Climatic conditions had a minor role on tree growth response to resin tapping. Our main conclusion is that the effect of resin extraction on growth is age-dependent. Our results encourage the co-production of resin and wood on maritime pine stands over 40 years old where resin extraction impact on growth is negligible.

**Keywords** Maritime pine · Resin tapping · Tree-ring width · Radial growth · Cambial age

## Introduction

Resin tapping and harvesting is a traditional forestry activity that provides economical profit to rural populations and other services like fire surveillance and sustainable forest management (Pereira 2015; Soliño et al. 2018). Currently, there is a new demand for natural resin (Génova et al. 2014) leading to a price increase and to the resurgence of the activity in places where it was almost abandoned, such as in the Iberian Peninsula (Girón, 2021).

Pine resin extraction has been usually done alongside timber production in the Mediterranean basin (Soliño

et al. 2018), especially in Portugal and Spain, where *Pinus pinaster* Aiton is the main species used for resin harvesting. Despite this, it has been hypothesized that secondary metabolism and growth may be competing for the available photosynthates (Züst & Agrawal 2017) and thus, repeated mechanical resin tapping could be stimulating resin biosynthesis at the expense of other carbon-dependent processes such as wood production (Du et al. 2021). Supporting this idea, lower growth rates and narrower tree-rings, as an estimate of biomass production, have been found in resin tapped pine stands (Papadopoulos 2013; Génova et al. 2014; Chen et al. 2015; Zeng et al. 2021). Papadopoulos (2013) showed a reduction of 14% in tree-ring width (6% of latewood) in tapped *Pinus halepensis* Mill trees, compared with non-tapped ones. Zeng et al. (2021) also found negative effects of resin harvesting on tree-ring width of *Pinus tabulaeformis* Carr, although it only persisted for two years after the extraction ended. But, positive (Tomusiak & Magnuszewski, 2013; van der Maaten et al. 2017) or no resin tapping effects (Rodríguez-García et al. 2015; Du et al. 2021) on tree-ring formation have also been reported, suggesting low

Communicated by Thomas Seifert.

✉ Mikael Moura  
mikael.moura@student.uc.pt

<sup>1</sup> Centre for Functional Ecology – Science for People & the Planet, Department of Life Sciences, University of Coimbra, Calçada Martim de Freitas, 3000–456 Coimbra, Portugal

carbon competition between growth and resin, which according to Hilty et al. (2021) can be considered as growth *sensu lato*—the combination of reversible and irreversible biomass or volume changes. Growth stimulation in tapped trees could also reflect reaction tissues if annual growth was measured too close to the tapping wound.

The contradictory results found in the literature concerning resin extraction effects on wood production might be also related to the studied species, the sampling design, and the resin tapping method and intensity. For example, Zeng et al. (2021) sampled both tapped and untapped sides to represent the overall tree-ring growth and observed that resin tapping in *P. tabulaeformis* promotes asymmetrical radial growth, with narrower rings near the tapped face, while Garcia-Forner et al. (2021a, b) showed the opposite pattern in *P. pinaster*. It was also observed that wood reaction to tapping decreases when getting away from the wound upward as shown in *Pinus sylvestris* L (Tomusiak & Magnuszewski, 2013). Thus, resin extraction impact on growth will depend on the sampling design performed (Williams 2017) hindering comparisons between studies. But even when the same sampling procedure and species are used, other variables, e.g., environmental conditions, can affect growth sensitivity to resin harvesting (Garcia-Forner et al. (2021a, b).

Tree growth and resin production are both driven by environmental conditions and depend on carbon availability to produce new cells, which include the resin duct complex formed by the resin canal and the adjacent resin-producing epithelial cells that synthesize the defensive terpenoid mixture. In *P. pinaster*, some studies connect favorable temperatures, solar radiation and potential evapotranspiration with increasing tree growth and resin duct traits or resin yield (Rodríguez-García et al. 2014, 2015), variables that are phenotypically and genetically correlated (Vázquez-González et al. 2019). However, literature is still scarce when it comes to understand the resin tapping impact on wood production under a climatic gradient, the key factor to identify the environmental conditions that maximize both resin and wood production. Another variable that should be considered is the species provenance, as growth, as well as resin yield are subjected to strong genetic control varying among families and provenance regions (Vázquez-González et al. 2021). If resin harvesting depletes carbon stocks, wood production in fast-growing populations could be more susceptible to resin extraction than in slow-growing ones (Zeng et al. 2021).

Stand and tree characteristics and forest management are also important factors that should be considered since they may determine resources availability at individual level, and thus affect their growth rate and production of defenses. For instance, at tree level, resin yield increases at lower stand densities (McDowell et al. 2007; Rodríguez-García et al. 2014, 2015; Zas et al. 2020). In high density stands, trees grow less, and smaller trees can be more affected (e.g., *P.*

*tabulaeformis* in He & Wang 2021) or end-up being less resilient to drought (Fernández-de-Uña et al. 2015; Bottero et al. 2017). The age of the trees affects their vitality, growth, and their resin yield too. In this regard, Zas et al. (2020) found that even if the mechanisms remain elusive, resin production in *P. pinaster* increases as trees get older. Considering that in woody plants photosynthesis tends to decrease with age (Yoder et al. 1994; Bond 2000), it is necessary to understand whether resin yield increases as age increases due to carbon allocation shifts from growth to defenses, or because of higher carbon reserves in older trees.

Our aim is to understand whether resin tapping affects tree-growth rates, and how tree, stand characteristics and edaphoclimatic conditions modulate it. We have studied six populations of *P. pinaster* along a latitudinal gradient in Portugal and assessed tree-ring width before and during resin tapping. Our hypotheses are that: i) resin tapping reduces tree-ring width due to shifts in carbon allocation to sustain resin production, but the impact will depend on ii) local conditions, and iii) the characteristics of the trees, i.e., trunk girth and height, tree density and competition between neighbors, with younger trees being more affected due to a lower capacity to acquire carbon and/or lower carbon reserves.

## Materials and methods

### Study sites and resin harvesting

Maritime pine (*Pinus pinaster* Aiton) is a fast-growing heliophytic tree species with a great ecological plasticity and varied climatic spectrum (Farjon 2010). In Portugal, this evergreen conifer represents 22% of the total forest area (ICNF 2015) and is the main species used for resin tapping. According to Portuguese law (see Art. 4<sup>th</sup> DL n.º 181/2015) resin tapping is performed in trees with more than 63 cm of trunk girth at 1.30 m from the ground, which is in line with previous studies that recommended resin tapping only in stands older than 30 years (Zas et al. 2020). Tapping starts on the trunk base at 20 cm above the ground by removing the bark and phloem every 15–20 days and continues upward for four years and up to 2 m high. Afterward, when tree size allows it, another vertical row is started 10 cm apart from the first wound and following the same procedure. Usually, the tapping season goes from March until November, coinciding with the radial growth of maritime pine.

For this study, we selected six stands of *P. pinaster* for resin production in the Northern and Central regions of Portugal, areas where this activity is traditionally done. Selection criteria of the stands was based on three main aspects: (1) to have a minimum of five years of tapping exploitation; (2) to have been exploited with only one resin wound

at a time and thus harmonizing the resin tapping intensity between stands; (3) to have used the same resin extraction methodology. We selected paired stands relatively close climatically, but different in the average age to assess its role on the impact of resin tapping on growth. The selected populations were located at Paredes de Coura (PCO), Vila Pouca de Aguiar (VPA), Olho (OLH), Soure (SOU), Marinha Grande (MGR) and Pataias (PAT) (Fig. 1). Across study sites, resin tapping occurred between 2013 and 2019 (Table 1). The number of tapped years was 5 in OLH and SOU; 6 in PCO, PAT and VPA and 7 in MGR (list of abbreviations in Table S1).


### Climatic data collection

Climatic data for each location was downloaded from the Climate Explorer of the Royal Netherlands Meteorological Institute (<https://climexp.knmi.nl>) using grid points 0.5 or

0.25. Due to their close proximity, MGR and PAT were located at the same grid point, and thus shared climatic data. Monthly time series of mean air temperatures (average, maximum and minimum; °C) and total precipitation (mm month<sup>-1</sup>) were obtained from E-OBS 21.0e 0.25. Self-calibrating Palmer Drought Severity Index (PDSI) calculated from Climatic Research Unit Time Series (CRU TS) data was obtained for 1901–2017 0.5° Global 3.26 early data set, and vapor pressure (hPa) from CRU TS 4.04 0.5° data set. Vapor pressure deficit (VPD, hPa) was calculated from average temperature and vapor pressure using the following formula:

$$VPD = 6.11 * \exp((17.27 * T) / (T + 273.3)) - VP \quad (1)$$

where *T* and *VP* are the means of the monthly average air temperature and vapor pressure, respectively. Maximum and minimum VPD were also calculated for each location using the respective temperatures and vapor pressure.



**Fig. 1** Location and climatic diagrams of the study sites. The color in the map represents the mean annual precipitation minus potential evapotranspiration (P-PET, in mm) for the period 1990–2019. Negative P-PET values indicate drier sites. For each site, an ombrothermic diagram is shown representing the average monthly air temperature

(*T*, in red) and precipitation (*P*, in blue) for the period 1990–2019, and the gray area indicates the drought period ( $2 T > P$ ). The annual average of air temperature and precipitation for a period of 30 years is displayed at the top of each climatic diagram

**Table 1** Stand, tree, and resin tapping characteristics for each site (PCO—Paredes de Coura; VPA—Vila Pouca de Aguiar; OLH—Olho; SOU—Soure; MGR—Marinha Grande; PAT—Pataias)

Location Sites		PCO	VPA	OLH	SOU	MGR	PAT	
Stand characteristics	Latitude	41.87	41.47	40.33	40.05	39.73	39.63	
	Longitude	−8.60	−7.52	−8.65	−8.59	−8.96	−8.96	
	Altitude (m a.s.l.)	479	791	71	49	100	98	
	Soil organic matter (%)	21.87±0.86 f	16.03±1.14 e	4.72±0.41 b	11.98±1.19 d	5.78±0.88 c	3.17±0.19 a	
	Soil pH	4.07±0.03 a	4.25±0.05 b	5.01±0.06 d	4.61±0.06 c	5.43±0.08 e	6.52±0.05 f	
	Tree density (trees/ha)	–	517±42 d	554±54 d	363±33 a	357±32 b	427±23 c	
Tree characteristics	Cambial age (years)	34±0.36 c	44±0.53 d	29±1.29 b	49±1.71 e	58±0.27 f	25±0.23 a	
	Trunk girth (cm)	98.0±3.5 b	88.1±3.0 a	102.9±2.1 c	96.0±4.0 c	108.8±1.6 d	108.4±1.5 d	
	Total height (m)	–	18.1±0.3 a	21.5±0.6 d	19.8±0.4 c	22.4±0.2 e	19±0.3 b	
	Trunk foliated length (m)	–	5.8±0.4 b	6.1±0.4 c	5.7±0.3 b	5.2±0.3 a	5.3±0.3 a	
	Neighbor trees (no.)	–	3.06±0.34	3.40±0.44	1.85±0.26	1.8±0.26	2.35±0.18	
	Neighbors' trunk girth (cm)	–	84.0±2.1 b	63.3±3.7 a	88.5±4.5 c	95.0±2.3 d	102.5±4.0 e	
	Height of neighbor trees (m)	–	18.2±0.3 b	15.1±0.7 a	17.7±0.5 c	22.1±0.2 d	17.8±0.5 e	
	CI <sub>TG</sub>	–	2.96±0.34 d	2.30±0.28 b	2.76±1.21 c	1.60±0.24 a	2.23±0.17 b	
	CI <sub>H</sub>	–	3.09±0.34 d	2.34±0.25 c	1.68±0.22 a	0.25 b	2.23±0.20 c	
	Resin Harvest	Tapping period (years)	2013–2018	2013–2018	2015–2019	2015–2019	2013–2019	2014–2019
		Number of tapped years	6	6	5	6	7	6
Tapped side (°)		–	184±36 ab	236±15 b	195±14 ab	154±18 ab	119±19 a	

CI<sub>TG</sub> and CI<sub>H</sub> are the two competition indexes using the trunk girths and heights of the neighbor trees divided by the trunk girths and heights of the sampled trees, respectively. Means and standard errors (±SE) are shown for variables measured at tree level. Different letters represent the statistical differences between sites for each parameter ( $p < 0.05$ )

## Stand and tree characteristics

Twenty *P. pinaster* trees per location were selected to investigate growth responses and the structure of the stand. We first evaluated the trees' canopy and then selected healthy trees with average trunk girth for each stand. Trunk girth at breast height (TG), total height and tree height until the first branches with green needles were measured. The difference between the two last variables was used as an estimate of the tree canopy size (trunk foliated length).

To quantify the degree of competition between trees and its potential effect on radial growth of selected trees, we calculated two competition indices based on trees' height and trunk girth (CI<sub>H</sub> and CI<sub>TG</sub>, respectively). This information is not available for PCO because trees were felled before collecting all data. We first identified all living trees in a circumference of 5-m radius from the trunk of each target tree, and then measured their trunk girth at breast height and the two described heights. The number of present neighbors was used to calculate tree density of the location (trees ha<sup>−1</sup>). CI<sub>H</sub> and CI<sub>TG</sub> were calculated as the sum of the ratio between each competitor tree in the circumference and the target tree as follows:

$$CI_{TG} = \sum tg_i / tg \quad (2)$$

where  $tg_i$  is the trunk girth at breast height of  $i^{\text{th}}$  competitor and  $tg$  is the trunk girth at breast height of the target tree; and

$$CI_H = \sum h_i / h \quad (3)$$

where  $h_i$  is the trees' height of  $i^{\text{th}}$  competitor and  $h$  is the height of the target tree.

Two soil samples per target tree were also collected at 50 cm (0–10 cm depth) from the north and south side of the trunk. Pooled samples were dried at 60 °C for four days and sifted with a 2-mm sieve. Soil pH was measured with a pH meter (JENWAY Model 3505) in a 1:2.5 (mass/volume, soil/water) suspension. The percentage of organic matter (OM) was assessed by loss on ignition at 450 °C for 4 h in a muffle furnace (Nelson & Sommers, 2018; Roberston 2011).


## Dendrochronological sampling and processing

Wood increment cores were collected on the 20 target trees per site in 2019 at PCO and VPA and in 2020 on the remaining sites. Two cores were taken from each tree at breast height (ca. 1.3 m), with an increment borer of 12-mm diameter. Core extraction was done between resin-tapping wounds, as all studied trees had been tapped for more than four consecutive years. The core between tapping wounds represents the resin tapping face and the one from the opposite side



represents the untapped face (Fig. 2). The orientation of each core was recorded to verify if the exposition of the tapping wound influenced growth response to tapping within and between sites.

After harvesting, each core was immersed in 99% acetone for 24 h at 4 °C to clean the samples from resin, dirt, and debris. Then, they were air dried, glued on wooden mounts and progressively polished (150, 400, 800 and 1200G) with a belt sander (Bosch GBS 100 AE). Polished samples were scanned at 2400 dpi and 48 bits in RGB using a photo-scanner (EPSON Perfection 4990 Photo) and the EPSON Scan software (2020 Seiko Epson Corporation). Increment core images were analyzed using the *xRing* package for R to automatically detect tree-ring borders and to measure tree-ring width (*TRW*) (Campelo et al. 2019; R Core Team 2020). After visually cross-dating by comparing *TRW* within tree (tapped and untapped sides) and site, the number of annual rings from the pith was used to determine the cambial age. The pith was not present in 1 tree at VPA, PCO, MGR and SOU and in 3 trees at OLH. Trees with correlations below 0.4 were discarded for further analysis. Final chronologies were computed with 15 trees for OLH, 18 for PCO and SOU, 20 for MGR, PAT and VPA. To remove size and age-related trends, a cubic smoothing spline was applied to each individual ring-width time-series. Due to the large variation in



**Fig. 2** Representation of an increment core sampled on the trunk of a tree with two adjacent tapping wounds. Concentric circles represent annual tree-rings. The green area represents the wound area where resin tapping—bark and phloem removal—was performed, and the red arrow shows the direction of the increment core, which starts between resin tapping wounds (tapped side) and ends on the opposite side (untapped side) of the trunk. The distance between tapped faces is around 10 cm according to the Portuguese law regulating the harvesting and marketing of pine resins (see Art. 4th DL n. ° 181/2015)

the average length of ring-width time series between sites, the recommendations of Klesse (2021) were followed to define the length of the spline. Thus, the grand mean of the mean of the time-series length for each site was used. In doing so, each ring-width time-series was detrended using a spline with a frequency response cut-off of 0.5 and a length of 26 years to enhance the climatic signal and to allow inter-site comparisons. Autoregressive modeling was performed on each standardized series to remove temporal autocorrelation and to maximize the climatic signal. For each site, the obtained residual ring-width indices (*RWI*) were averaged using a bi-weighted mean to produce a site chronology. For the common interval to all sites, the statistical quality of each chronology was evaluated using the following variables: the expressed population signal (*EPS*) that indicates the degree to which a given chronology approaches an infinitely replicated chronology (Wigley et al. 1984a, b), the *r.tot* which is the mean of all correlation between different cores, *r.wt* which is the mean of correlations between series from the same tree over all trees and *r.bt* which is the mean inter-series correlation between all series from different trees. The chronologies and detrending were developed using *dplR* (Bunn 2010; R Core Team 2020) and *detrendR* (Campelo et al. 2012) packages for R version 3.6.2 (R Core Team 2020).

## Statistics

We characterized the climatic differences between sites by conducting a principal component analysis (PCA) of the main climatic variables between 1999 and 2019, the common period for all the chronologies. Variables with low contribution percentage among principal components were removed from the PCA. The variables included in the final PCA were: *mat* (mean temperature), *pp* (mean annual precipitation), *ts* (temperature seasonality), *maxvpd* (maximum vapor pressure deficit) and *mpdsi* (mean Palmer Drought Severity Index). All variables were previously scaled, and the analysis was computed using the function *PCA* from the *FactoMineR* package (Lê et al. 2008) for R version 3.6.2 (R Core Team 2020).

The parameters of the studied trees (*TG*, tree height, trunk foliate length) and the structure of the stand (tree density, core exposition, soil *pH* and *OM*,  $CI_{TG}$  and  $CI_H$ ) between sites were compared using linear models. All response variables were normally distributed, except soil *pH* and *OM* that were square root transformed to meet the assumptions of parametric analyses. The respective tree/stand parameters were the dependent variable, site the fixed factor and tree the random factor. Bonferroni method was used for the pairwise post hoc tests and  $\alpha < 0.05$  as a significant level. These analyses were performed using the function from the *nlme* package v.3.1–14 (Pinheiro et al. 2020) and the *cld* function

from the *multcomp* during pairwise post hoc tests (Hothorn et al. 2008) function for R (R Core Team 2020).

To comprehend if resin tapping affects tree-ring width at site level, we compared the *TRW* during the tapping years and the same number of years before the start of resin harvesting, e.g., 2006–2012 before tapping vs. 2013–2019 during tapping in MGR. Then, linear mixed models were performed using *TRW* as dependent variables, square root transformed to meet a normal distribution. Site, tapping (before vs. during), and trunk side (tapped vs. untapped) were considered as fixed factors, while tree within site was considered a random factor. Pairwise post hoc tests with Bonferroni corrections were carried out to compare differences between trunk side, and period of tapping for each site. The same procedure was applied to *RWI* to assess the results after removing age influence on growth trends.

To compare resin harvesting impact between years and stands with different growth rates, we standardized data calculating the resin tapping impact (*RTI*), i.e., the ratio between *TRW* of each tapping year and the mean *TRW* for the 5 years before tapping started. The *RTI* corresponds to the resistance index proposed by Lloret et al. (2011), which estimates the performance during and before a disturbance, i.e., resin tapping in this study. We have used 5 years averages for the pre-disturbance period to reduce the weight of any natural trend, and to account for inter-annual climatic variability. Values > 1 and < 1 indicate that wider and narrower rings were produced during tapping relative to the previous period, respectively. To compare the *RTI* between tapping years and sites, we used linear mixed models where *RTI* was the dependent variable, site, and the tapping year the fixed factors and tree the random factor. More complex models with autocorrelation were considered but did not improve model performance according to Akaike and Bayesian information criteria. When significant effects were detected, a pairwise post hoc test with Bonferroni corrections was applied to account for multiple comparisons between years and sites. For these analyses, it was used the *lme* function from the *nlme* package v.3.1–14 (Pinheiro et al. 2020) and the *cld* function from the *multcomp* during pairwise post hoc tests (Hothorn et al. 2008).

To assess the effect of tree traits, stand characteristics, and climate on *RTI*, the Spearman's rank correlation coefficient was calculated within and across sites. The parameters used for the tree and stand characteristics were *TRW*, *TG*, tree height, tree foliated height, tree density, the competition indices ( $CI_{TG}$  and  $CI_H$ ), soil *pH*, soil *OM* and cambial age. For the PCO site, where trees were felled before we could collect data for radial growth. Only *TRW*, *TG*, and the soil *pH* and *OM* variables were used. The climatic parameters used were based on the scores of the first, second, third, and fourth principal components of the PCA (*mat*, *ts*, *pp*, *maxvpd* and *mpdsi*). These analyses

were conducted using *rcorr.adjust* function of the *Hmisc* package for R version 3.6.2 (R Core Team 2020).

## Results


### Climate and stand characteristics

The sampling sites showed a latitudinal climatic gradient with the coldest location, VPA, with an average temperature of 12.1 °C, also located at the highest altitude, 791 m (Table 1, Abbreviations in Table S1). The warmest location was SOU, whereas MGR and PAT were the driest ones, with negative P–PET values (Fig. 1). PCO showed the highest precipitation levels (1471 mm on average) and highest average *VPD* (6.88 hPa) (Fig. 1, Table S2). The average Palmer Drought Severity Index (*PDSI*) was close to 0 and similar between sites, but slightly more negative in the lower latitudes (Table S2).

The PCA retained 5 climatic variables: *mat*, *pp*, *ts*, *maxvpd* and *mpdsi*. The two first principal components (henceforth described as PC1 and PC2) explained 38.9% and 23.2% of the overall variance, respectively (Fig. 3). The variables *ts* and *maxvpd* were positively related to PC1, while *mat*, *pp* and *mpdsi* were negatively correlated. Precipitation (*pp*) was negatively correlated with PC2, while the other variables were positively related. The PC1 and PC2 aggregated the sites in three main groups: i) PCO, ii) VPA and iii) OLH, SOU, MGR and PAT (Fig. 3). PCO was positively related to *pp* and *mpdsi*, and negatively to *mat*, while VPA had a positive correlation with *ts*. The rest of the locations correlated positively with PC2, with *mat*, *maxvpd* and *mpdsi* (Fig. 3).

Soil *OM* and *pH* were different among sites (Table 1) with PCO and VPA (Table 1) showing higher *OM* and lower *pH* values. Across sites, average cambial age varied from 58 to 29 years-old *TG* and height between 88 and 108 cm and 18 and 22 m, respectively (Table 1). The biggest and tallest trees were found in MGR, the oldest stand; however, *TG* in PAT, the stand located at the same climatic grid point, was similar, even if the latter was the youngest sampled population (Table 1). Marinha Grande (MGR) and SOU showed, on average, less than two neighbors, in contrast with the 3.4 per target tree identified in OLH; however, competitor trees were smaller in the latter (Table 1). Olho (OLH) was the densest stand with the highest inter-tree competition (554 trees/ha and  $CI_{TG}$  2.30), and MGR the one with the lowest density and competition indices (357 trees/ha and  $CI_{TG}$  1.60). Across locations, the tapped side of the trunk (tapped increment core) tended to be exposed to south, varying between southeast and southwest (Table 1).

**Fig. 3** Principal component analysis of the climate data [annual mean temperature (*mat*), annual mean precipitation (*pp*), temperature seasonality (*ts*), maximum vapor pressure deficit (*maxvpd*) and mean Palmer Drought Severity Index (*mpdsi*)] for the six study sites [Paredes de Coura (PCO), Vila Pouca de Aguiar (VPA), Olho (OLH), Soure (SOU), Marinha Grande (MGR) e Pataias (PAT)] with the two principal components explaining 62.1% variance overall. Different sites are depicted by different symbols and colors. Due to its proximity, MGR and PAT share the same climatic data




**Tree-ring width before and during tapping**

In general, the tree-ring width index series considering the common interval within (Table S3) and between sites (Table S4; 1999–2019) showed *EPS* values above 0.85, except in OLH and SOU that showed slightly lower values from 1999 to 2019 and including the tapped and untapped side of the trunk. PAT, the youngest stand, was the site with the highest average *TRW* values (5.53 mm), followed

by OLH (4.22 mm), PCO (3.73 mm), VPA (2.54 mm), MGR (2.27 mm) and SOU (1.91 mm). In general, all chronologies showed the age growth trend producing narrower tree-rings with increasing tree age (Fig. 4). *TRW* of the untapped side was smaller in PCO and OLH; however, this trend was present before the resin tapped period (Fig. 4). To understand if *TRW* was affected by resin harvesting, we compared the mean *TRW* between the resin tapped years and the same number of years before tapping

**Fig. 4** Average tree-ring widths (*TRW*, mm) between 1999 and 2019 for the study sites: Paredes de Coura, (PCO), Vila Pouca de Aguiar (VPA), Olho (OLH), Soure (SOU), Marinha Grande (MGR) and Pataias (PAT). Red and blue lines represent the untapped and tapped side of the trunk, respectively. The gray shadow represents the resin tapping period. The equivalent number of years before the beginning of tapping for each site is shown in dark yellow. Resin extraction started in 2013 in PCO, VPA and MGR, in 2014 in PAT, and in 2015 in OLH and SOU. Note that *TRW* scale varies between plots




calculated separately for the tapped and untapped trunk sides (Fig. 5). Pine trees in two of the youngest populations, PCO and OLH, showed a significant decrease of the *TRW* during the tapping period on both sides of the trees, while PAT—the youngest and fastest growing

population—showed a *TRW* decrease only in the untapped side (Fig. 5). All the other sites (VPA, SOU and MGR) showed no significant growth decrease during tapping, regardless of the side of the tree (Fig. 5). In MGR, the stand with the oldest tree population, *TRW* increased



**Fig. 5** Boxplots of the tree-ring widths (*TRW*, mm) for each side of the tree (tapped in blue and untapped in red) before and during resin tapping. Different letters indicate statistically significant differences between groups at each location ( $p < 0.05$ )



during the tapping period on the tapped side of the trunk (Fig. 5). The same analysis was performed using the tree-ring-width index (*RWI*), where the *TRW* age trend was removed (Fig. S1). In this case, *RWI* increased during the tapping period in VPA and MGR, regardless of the side of the tree (Fig. S1). OLH showed the opposite trend, that

is, *RWI* was higher before than after the tapping started, while PCO, PAT and SOU showed no significant differences (Fig. S1).


## Resin tapping impact

On average, the *RTI* was  $< 1$  in the youngest populations (PCO = 0.72, OLH = 0.84 and PAT = 0.85) and  $> 1$  in the oldest populations (VPA = 1.17, SOU = 1.18 and MGR = 1.41) (Fig. 6). Despite *RTI* fluctuations between different tapping years, no clear pattern associated with


cumulative stress due to repeated resin harvesting was detected (Fig. 6).

The Spearman's rank correlation coefficients between *RTI*, tree, and stand characteristics across sites were significant and positive for *TRW* (+0.38), cambial age (+0.53), total height (+0.24), and soil *OM* (+0.11), and negative with *CI<sub>TC</sub>* (-0.22) and soil pH (-0.14) (Table S5, Fig. 7).

**Fig. 6** Resin tapping impact (*RTI*) per site during the years of resin tapping. Values of *RTI*  $< 1$  and  $> 1$  indicate narrower and larger tree-rings during resin extraction, respectively, in comparison with pre-disturbance. Letters denote statistical differences between resin tapping years within site at  $p < 0.05$ . Sites are organized from the youngest to the oldest stands (from left to right). Resin extraction started in 2013 in PCO, VPA and MGR, in 2014 in PAT and in 2015 in OLH and SOU



**Fig. 7** Spearman's rank correlations coefficients between *RTI* with cambial age **a**, total height **b**, trunk girth **c** and  $CI_{TG}$  **d** across sites. Each gray dot represents individual trees (average *RTI* during the tapping period and the characterization variable measured during the sampling campaign) and the blue dots represent the average value per site with its respective standard error bar (SE) and regression line (in red). Relationships between variables are represented by dashed lines when non-significant and by solid lines when significant. The Spearman's correlation coefficient ( $R$ ) and the  $p$ -values are shown (for  $p < 0.05$ )




Within sites, *RTI* was positively correlated with *TRW* in all cases (Table S5). In general, *TG* was strongly and positively associated with *RTI* in VPA and PAT, to total height in PAT and OLH, but to soil *OM* only in PCO. Negative *RTI* were shown for cambial age in MGR and PAT (Table S5).

Spearman correlations were carried out between the *RTI* and the scores of the first, second, third and fourth

components of the PCA (Fig. 8). The scores of the principal components were not significantly associated with *RTI*, except for the PC4 that preserved only the 10.6% of the climatic variance and was mainly explained by *mat* (32.8%). In the latter, with the increasing *mat*, the *RTI* decreases to values below 1 (Fig. 8d).

**Fig. 8** Spearman's rank correlation coefficients between the resin tapping impact (*RTI*) and the scores of the first **a**, second **b**, third **c** and fourth **d** climatic principal components (PC1, PC2, PC3 and PC4; see Fig. 3) for all studied locations. Each dot represents each tapping year for each site. Relationships between variables are represented by dashed lines when non-significant ( $p < 0.05$ ) and by solid lines when significant ( $p < 0.05$ ). The PC1 is more weighted by the variable *ts* and the PC2 by *pp*. PC3 has *mpdsi* has the highest contributed variable and PC4 is with *mat*. The Spearman correlation coefficient (*R*) and the *p*-values are shown



## Discussion

The aim of this study was to understand if resin tapping affects wood growth of *P. pinaster*, and whether tree, stand characteristics and edaphoclimatic conditions can modulate it or not. Our results show that radial growth is not systematically reduced on maritime pine trees exploited

for resin production and hint that negative impacts are more frequent in younger (< 30 years-old) than older populations as hypothesized. This result indicates that, as well as being a profitable forest activity, resin and timber co-production in maritime pine may be sustainable, especially when performed in populations older than 40-year-old.



In general, when the *TRW* before and during resin tapping is compared, the youngest populations (PCO, OLH and PAT) showed a decreasing growth trend during resin extraction, while the oldest populations showed no significant differences (VPA, SOU) or even wider rings during this period (MGR). Tomusiak & Magnuszewski (2013) and van der Maaten et al. (2017) also showed positive influence of tapping on radial growth on Scots pine and no systemic reductions in growth as we found in the older stands. On the contrary, Chen et al. (2015) showed that younger tapped Masson pine trees were able to enlarge and repair their trunks faster than older trees, as their tree-ring index only declined by 4.8%. However, they only investigated two sites, and no details about stand structure were given. Discrepancies can also be partially explained by the use of non-detrended series. Studying resin tapping impact through tree-ring width may be confusing because the biological trend associated with tree age is still present. Yet, our results consistently showed that growth resistance increases in older stands when *RWI* instead of the *TRW* was considered. In this case, two of the oldest stands (VPA and MGR) showed higher growth indices during tapping than the pre-disturbance period. Although resin production was not measured here, Zas et al. (2020) showed that resin production increases considerably with tree age in maritime pine stands growing under similar climatic conditions in northern Portugal. Determining a minimum age to start resin harvesting in this species becomes, therefore, a key factor to design sustainable timber and resin forest exploitations, and future studies should shed light on this.

Besides the stand age, other parameters must be taken into account when trying to understand growth sensitivity to resin harvesting, for instance the design of the increment cores' sampling. Wood formation in the trunk of tapped pines is not uniform as growth is suppressed in the region due to the mechanical injury of the bark and phloem tissues, and asymmetrical rings can be produced in the remaining surface depending on their distance to the wound (Grissino-Meyer et al. 2001). Although our study did not measure the effect of tapping on wood properties, other studies have shown that resin tapping results in higher wood deformation and elasticity (Silva et al. 2018), but also increases wood density (especially on the tapped side), which modifies its mechanical properties (García-Iruela et al. 2016; Silva et al. 2018; Wu et al., 2022). Furthermore, if the wood core is extracted too close to the wound, the increase in *TRW* can just be related to the wood reaction to the injury. To take this into account, Zeng et al. (2021) extracted six increment cores per tree, and Papadopoulos (2013) measured *TRW* and latewood in three radial directions in Aleppo pines. Nonetheless, impact assessment, as well as comparisons between studies, is usually difficult as most of the sampling designs usually include the extraction of a single increment core per tree and

height taken from the tapped or untapped face (Tomusiak & Magnuszewski., 2013; Hood & Sala 2015; Zas et al. 2020; Du et al. 2021). In this study, we harvested stands subjected to the same tapping intensity (number of wounds at a time and tapping frequency), and collected wood cores using the same methodology, so that the comparison between sites can disclose the effect of stand age and edaphoclimatic characteristics among the studied sites.

Across sites, the locations characterized by taller trees producing wider tree-rings, low inter-tree competition at *TG* level, and greater content in soil organic matter tended to show higher growth rates during tapping than the previous period (*RTIs* above 1). On the one hand, larger and taller trees may have better access to light and can potentially photosynthesize more, acquiring more carbon that can be used for either resin or wood production. Furthermore, taller trees usually have deeper roots and are less sensitive to variations in precipitation because they can tap deeper water sources and thus, keep high transpiration and photosynthesis rates for longer under drought (Brando 2018; Giardina et al. 2018). On the other hand, the negative  $CI_{TG}$  correlation with *RTI* suggests that competition for resources could affect the capacity to respond to harvesting, as well as shown in response to other disturbances (Rozendaal et al. 2020; Marqués et al. 2021). In addition, it has been shown that at least in young *Pinus oocarpa* Schiede ex Schltdl, high competition levels lead to decreasing resin yield, although the opposite can occur in adult trees (Egloff 2020). Therefore, and specially in young stands, it is necessary to carry out forest management practices such as thinning to reduce inter-tree competition, as well as the associated growth and resin production constraints.

Within sites, only PCO—the tree population growing in more acidic soils—showed significant correlations between soil organic matter, *pH* and *RTI*, positive and negative, respectively. Organic matter is an important source of nitrogen, a macronutrient for plants and central for photosynthesis, with leaves with higher amounts of nitrogen showing a higher photosynthetic capacity and increasing the growth potential of the trees. The negative association between soil *pH* and *RTI* could simply reflect the effect of this variable on the mineralization of the organic matter, and/or to a higher nutrient availability under more acidic soils (Curtin et al. 1998).

Selected pine stands in this study could represent different *P. pinaster* provenances (ICNF 2012) and thus, their annual growth could be strongly influenced by climatic conditions and, to a lesser extent, by genetic plasticity (Rozas et al. 2020). Despite this, the climatic sensitivity of tapped and untapped trees appears to be similar, as shown by van der Maaten et al. (2017). In their study, tapped and non-tapped *P. sylvestris* trees presented comparable sensitivities even in the occurrence of an extreme

climatic event. In line with this, our results showed that the climatic conditions had a marginal effect on *RTI* across stands. Only 10% of *RTI* variance was explained by PC4 indicating that growth sensitivity across sites increases with increasing average temperatures. Without taking into account the potential effect of temperature on resin production and considering the studied region—where resin is traditionally harvested—warmer places seem to be less profitable for timber production when the activity is combined with resin harvesting, a trend that could be exacerbated by rising temperatures.

In conclusion, this study showed that the age of maritime pine stands had a strong effect on the impact of resin extraction on wood growth, with younger stands (< 30 years-old) showing a tendency to form narrower rings—reduced wood production, and a lower capacity to buffer the resin extraction disturbance compared with older stands. Nonetheless, we did not have access to the resin yield of the different stands, an important information to deepen our knowledge on the impact of resin extraction on the carbon economy of maritime pine. Based on our results, to maximize resin and wood co-production, resin extraction should be performed in populations with more than 40 years of age, and on managed stands with controlled tree density to reduce competition and thus provide greater growth resistance to resin extraction.

**Supplementary Information** The online version contains supplementary material available at <https://doi.org/10.1007/s10342-023-01568-7>.

**Acknowledgements** We are very grateful for the help provided by: RESIPINUS for providing the contacts of the resin tapping operators and landowners; Gestão Integrada e Fomento Florestal Lda. as well as J. Lousada e E. Silva for their collaboration during the samplings in PCO and VPA; Pedrosa e Irmãos, Lda. and Mr. Carlos Simões for helping us in SOU; Mr. Manuel Santos, resin worker at OIH for his advice and permission to sample; and Mr. Luís Carpalhoso and Costa e Irmãos Lda for allowing us to conduct this study at PAT and MGR. The first author wishes to thank Ana Carvalho for her help and support on the field and laboratory.

**Author contributions** All authors contributed to the study conception and design. Field samplings, data collection and analysis were performed by MM, NG-F and FC. All authors contributed to interpret the data. MM wrote the manuscript with the help of the other authors, and all read and approved the final version of the manuscript.

**Funding** Open access funding provided by FCT/IFCCN (b-on). This study was co-financed by the Portuguese Foundation for Science and Technology (FCT) and the European Regional Development Fund through the project PTDC/ASP-SIL/31231/2017. The corresponding author is funded by a PhD Student Grant (SFRH/BD/145914/2019) from the FCT. The investigation was carried out at the R&D Unit designated Center for Functional Ecology—Science for People and the Planet (project UIDB/04005/2020) supported by Portuguese national funds.

**Data availability** Data will be made available upon request.

**Code availability** Code will be made available upon request.

## Declarations

**Conflict of interest** The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Bond BJ (2000) Age-related changes in photosynthesis of woody plants. *Trends Plant Sci* 5(8):349–353. [https://doi.org/10.1016/S1360-1385\(00\)01691-5](https://doi.org/10.1016/S1360-1385(00)01691-5)
- Bottero A, D'Amato AW, Palik BJ, Bradford JB, Fraver S, Battaglia MA, Asherin LA (2017) Density-dependent vulnerability of forest ecosystems to drought. *J Appl Ecol* 54(6):1605–1614. <https://doi.org/10.1111/1365-2664.12847>
- Brando P (2018) Tree height matters. *Nat Geosci* 11(6):390–391. <https://doi.org/10.1038/s41561-018-0147-z>
- Bunn AG (2010) Statistical and visual crossdating in R using the dplR library. *Dendrochronologia* 28(4):251–258. <https://doi.org/10.1016/j.dendro.2009.12.001>
- Campelo F, García-González I, Nabais C (2012) detrendeR: a Graphical User Interface to process and visualize tree-ring data using R. *Dendrochronologia* 30(1):57–60. <https://doi.org/10.1016/j.dendro.2011.01.010>
- Campelo F, Mayer K, Grabner M (2019) xRing—An R package to identify and measure tree-ring features using X-ray microdensity profiles. *Dendrochronologia* 53:17–21. <https://doi.org/10.1016/j.dendro.2018.11.002>
- Chen F, Yuan YJ, Yu SL, Zhang TW (2015) Influence of climate warming and resin collection on the growth of Masson pine (*Pinus massoniana*) in a subtropical forest, southern China. *Trees – Struct Funct* 29(5):1423–1430. <https://doi.org/10.1007/s00468-015-1222-3>
- Curtin D, Campbell CA, Jalil A (1998) Effects of acidity on mineralization: pH-dependence of organic matter mineralization in weakly acidic soils. *Soil Biol Biochem* 30(1):57–64. [https://doi.org/10.1016/S0038-0717\(97\)00094-1](https://doi.org/10.1016/S0038-0717(97)00094-1)
- Du B, Luan Q, Ni Z, Sun H, Jiang J (2021) Radial growth and non-structural carbohydrate partitioning response to resin tapping of slash pine (*Pinus elliottii* Engelm. var. *elliottii*). *J Forestry Research* 33(2):423–433. <https://doi.org/10.1007/s11676-021-01357-1>
- Egloff, P. (2020). Tapping *Pinus oocarpa* Assessing drivers of resin yield in natural stands of *Pinus oocarpa* Master thesis Philipp Egloff AV2019–25. March.
- Farjon A (2010) A handbook of the world's conifers. BRILL, Leiden, Boston, pp 533–1111
- Fernández-de-Uña L, Cañellas I, Gea-Izquierdo G (2015) Stand competition determines how different tree species will cope with a warming climate. *PLoS ONE* 10(3):1–19. <https://doi.org/10.1371/journal.pone.0122255>

- García-Fórner N, Campelo F, Carvalho A, Vieira J, Rodríguez-Pereiras A, Ribeiro M, Salgueiro A, Silva ME, Louzada JL (2021a) Growth-defence trade-offs in tapped pines on anatomical and resin production. *Forest Ecol Manag* 496:119406. <https://doi.org/10.1016/j.foreco.2021.119406>
- García-Fórner N, Carvalho A, Campelo F (2021b) Water and stem girdling affect the tracheids' number more than their shape in *Pinus pinaster* saplings. *Trees* 35:1921–1931. <https://doi.org/10.1007/s00468-021-02160-5>
- García-Iruela A, Esteban L, De Palacios P, García-Fernández F, De Miguel A, Vázquez E, Simón C (2016) Resinous Wood of *Pinus pinaster* Ait. *Physio-Mech Prop Bioresour* 11:5230–5241. [https://doi.org/10.1007/978-3-319-95432-5\\_5](https://doi.org/10.1007/978-3-319-95432-5_5)
- Génova M, Caminero L, Dochao J (2014) Resin tapping in *Pinus pinaster*: Effects on growth and response function to climate. *Eur J Forest Res* 133(2):323–333. <https://doi.org/10.1007/s10342-013-0764-4>
- Giardina F, Konings AG, Kennedy D, Alemohammad SH, Oliveira RS, Uriarte M, Gentine P (2018) Tall Amazonian forests are less sensitive to precipitation variability. *Nat Geosci* 11(6):405–409. <https://doi.org/10.1038/s41561-018-0133-5>
- Susana Girón. (2021). Spain's untapped "liquid gold." <https://www.bbc.com/travel/article/20211014-spains-untapped-liquid-gold>
- Grissino-Meyer HD, Blount HC, Miller AC (2001) Tree-ring dating and the ethnohistory of the naval stores industry in southern Georgia. *Tree-Ring Res* 57(1):3–13
- Hao-ran Wu, Yan-ru F, Xiao-yun N, Qi-fu L, Yan-jie Li, Jing-min J, Jian-er J (2022) Effects of resin-tapping year on wood properties of living *Pinus elliottii*. *Forest Res* 35(1):31–39. <https://doi.org/10.13275/j.cnki.lykxyj.2022.01.004>
- He R, Wang X (2021) Impact of Competition on the Growth of *Pinus Tabulaeformis* in Response to Climate on the Loess Plateau of China. *Plant Ecol* 223:353–368
- Hilty J, Muller B, Pantin F, Leuzinger S (2021) Plant growth: The what, the how, and the why. *New Phytologist* 232(1):25–41. <https://doi.org/10.1111/nph.17610>
- Hood S, Sala A (2015) Ponderosa pine resin defenses and growth: metrics matter. *Tree Physiol* 35(11):1223–1235. <https://doi.org/10.1093/treephys/tpv098>
- Hothorn T, Bretz F, Westfall P (2008) Simultaneous inference in general parametric models. *Biom J* 50(3):346–363. <https://doi.org/10.1002/bimj.200810425>
- ICNF. (2012). Regiões de proveniência de Portugal. Instituto Da Conservação Da Natureza e Das Florestas, Lisboa., 1–88. <https://www.icnf.pt/api/file/doc/7eba73fd7d9976f1>
- ICNF. (2015). 6.º Inventário Florestal Nacional (IFN6) - Relatório Final. 284. <http://www2.icnf.pt/portal/florestas/ifn/ifn6>
- Klesse S (2021) Critical note on the application of the "two-third" spline. *Dendrochronologia* 65:125786. <https://doi.org/10.1016/j.dendro.2020.125786>
- Lê S, Josse J, Husson F (2008) FactoMineR: an R package for multivariate analysis. *J Stat Softw* 25:1–18
- Lloret F, Keeling EG, Sala A (2011) Components of tree resilience: Effects of successive low-growth episodes in old ponderosa pine forests. *Oikos* 120(12):1909–1920. <https://doi.org/10.1111/j.1600-0706.2011.19372.x>
- Marqués L, Camarero JJ, Zavala MA, Stoffel M, Ballesteros-Cánovas JA, Sancho-García C, Madrigal-González J (2021) Evaluating tree-to-tree competition during stand development in a relict Scots pine forest: how much does climate matter? *Trees – Struct Funct* 35(4):1207–1219. <https://doi.org/10.1007/s00468-021-02109-8>
- McDowell NG, Adams GD, Bailey JD, Kolb TE (2007) The role of stand density on growth efficiency, leaf area index, and resin flow in southwestern ponderosa pine forests. *Can J for Res* 37(2):343–355. <https://doi.org/10.1139/X06-233>
- Nelson, D. W., & Sommers, L. E. (2018). Total carbon organic carbon and organic matter. *Methods of Soil Analysis: Part 3 Chemical Methods*, 5 7 961–1010 doi <https://doi.org/10.2136/sssabookse-r5.3.c34>
- Papadopoulos AM (2013) Resin tapping history of an Aleppo Pine forest in central Greece. *The Open Forest Science Journal* 6(1):50–53. <https://doi.org/10.2174/1874398601306010050>
- Pereira, J. M. R. (2015). Estimativa do potencial produtivo de resina em pinheiro-bravo no concelho de Castro Daire. Mestre em Engenharia Florestal e dos Recursos Naturais.pp 62.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Team, R. C. (2020). nlme: Linear and Nonlinear Mixed Effects Models [WWW Document]. R Packag. Version 3.1–151.
- R Core Team (2020) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>.
- Roberston, S. (2011). Direct estimation of organic matter by loss on ignition: methods. SFU Soil Science Lab (2011).
- Rodríguez-García A, López R, Martín JA, Pinillos F, Gil L (2014) Resin yield in *Pinus pinaster* is related to tree dendrometry, stand density and tapping-induced systemic changes in xylem anatomy. *For Ecol Manage* 313:47–54. <https://doi.org/10.1016/j.foreco.2013.10.038>
- Rodríguez-García A, Martín JA, López R, Mutke S, Pinillos F, Gil L (2015) Influence of climate variables on resin yield and secretory structures in tapped *Pinus pinaster* Ait. in central Spain. *Agric for Meteorol* 202:83–93. <https://doi.org/10.1016/j.agrformet.2014.11.023>
- Rozas V, Sampedro L, Vázquez-González C, Solla A, Vivas M, Lombardero MJ, Zas R (2020) Site conditions exert more control than genetic differentiation on modulation of secondary growth and climate sensitivity of *Pinus pinaster*. *Dendrochronologia* 63:125732. <https://doi.org/10.1016/j.dendro.2020.125732>
- Rozendaal DMA, Phillips OL, Lewis SL, Affum-Baffoe K, Alvarez-Davila E, Andrade A, Aragão LEOC, Araujo-Murakami A, Baker TR, Bánki O, Brienen RJW, Camargo JLC, Comiskey JA, Djuikouo Kamdem MN, Fauset S, Feldpausch TR, Killeen TJ, Laurance WF, Laurance SGW, Vanderwel MC (2020) Competition influences tree growth, but not mortality, across environmental gradients in Amazonia and tropical Africa. *Ecology* 101(7):1–11. <https://doi.org/10.1002/ecy.3052>
- Silva ME, Loureiro C, Gaspar MJ, Pires J, Ribeiro M, Loureiro C, Coutinho JP, Santos E, Carvalho A, Brito JL, Salgueiro A, Louzada JL, 2018. RESIMPROVE - Desenvolvimento de processos de produção e extração de resina de pinheiro para a melhoria da eficiência, racionalização e expansão da atividade. ISBN 978-989-704-264-5
- Soliño M, Yu T, Alía R, Auñón F, Bravo-Oviedo A, Chambel MR, de Miguel J, del Río M, Justes A, Martínez-Jauregui M, Montero G, Mutke S, Ruiz-Peinado R, García del Barrio JM (2018) Resin-tapped pine forests in Spain: Ecological diversity and economic valuation. *Sci Total Environ* 625:1146–1155. <https://doi.org/10.1016/j.scitotenv.2018.01.027>
- Tomusiak R, Magnuszewski M (2013) Effect of resin tapping on radial increments of Scots pine (*Pinus sylvestris* L.). *Lesne Prace Badawcze* 74(3):273–280
- van der Maaten E, Mehl A, Wilmking M, van der Maaten-Theunissen M (2017) Tapping the tree-ring archive for studying effects of resin extraction on the growth and climate sensitivity of Scots pine. *Forest Ecosyst* 4:1–7. <https://doi.org/10.1186/s40663-017-0096-9>
- Vázquez-González C, López-Goldar X, Zas R, Sampedro L (2019) Neutral and climate-driven adaptive processes contribute to explain population variation in resin duct traits in a mediterranean pine species. *Front Plant Sci* 10:1–12. <https://doi.org/10.3389/fpls.2019.01613>

- Vázquez-González C, López-Goldar X, Alía R, Bustingorri G, Lario FJ, Lema M, de la Mata R, Sampedro L, Touza R, Zas R (2021) Genetic variation in resin yield and covariation with tree growth in maritime pine. *Forest Ecol Manag* 482:118843. <https://doi.org/10.1016/j.foreco.2020.118843>
- Wang Z, Calderon MM, Carandang MG (2006) Effects of resin tapping on optimal rotation age of pine plantation. *J for Econ* 11(4):245–260. <https://doi.org/10.1016/j.jfe.2005.10.001>
- Wigley TML, Briffa KR, P. D. J. (1984a) On the average value of correlated times series, with applications in dendroclimatology and hydrometeorology. *J Clim Appl Meteorol* 23(1984):201–213
- Wigley TML, Briffa KR, Jones PD (1984b) On the average value of correlated time series, with applications in Dendroclimatology and Hydrometeorology. *J Climate Appl Meteorol* 23:201–213
- Williams R (2017) The effects of resin tapping on the radial growth of masson pine trees in South China-A case study. *Agricult Res Technol* 8(2):8–11. <https://doi.org/10.19080/artoaj.2017.08.555732>
- Yoder BJ, Ryan MG, Waring RH, Schoettle AW, Kaufmann MR (1994) Evidence of reduced photosynthetic rates in old trees. *Forest Sci* 40(3):513–527. <https://doi.org/10.1093/forestscience/40.3.513>
- Zas R, Quiroga R, Touza R, Vázquez-González C, Sampedro L, Lema M (2020) Resin tapping potential of Atlantic maritime pine forests depends on tree age and timing of tapping. *Industrial Crops and Prod* 157:112940. <https://doi.org/10.1016/j.indcrop.2020.112940>
- Zeng X, Sun S, Wang Y, Chang Y, Tao X, Hou M, Wang W, Liu X, Zhang L (2021) Does resin tapping affect the tree-ring growth and climate sensitivity of the Chinese pine (*Pinus tabulaeformis*) in the Loess Plateau, China? *Dendrochronologia* 65:125800. <https://doi.org/10.1016/j.dendro.2020.125800>
- Züst T, Agrawal AA (2017) Trade-offs between plant growth and defense against insect herbivory: an emerging mechanistic synthesis. *Annu Rev Plant Biol* 68(1):513–534. <https://doi.org/10.1146/annurev-arplant-042916-040856>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



## Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

[onlineservice@springernature.com](mailto:onlineservice@springernature.com)